

O-bearing Molecules in Carbon-rich Proto-Planetary Objects¹

F. Herpin², J. Cernicharo

Depto Física Molecular, I.E.M., C.S.I.C, Serrano 121, E-28006 Madrid, Spain

Received ** 1999; accepted ** 1999

¹ Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands and the United Kingdom) and with participation of ISAS and NASA.

²herpin@isis.iem.csic.es

ABSTRACT

We present ISO LWS observations of the proto-planetary nebula CRL 618, a star evolving very fast to the planetary nebula stage. In addition to the lines of ^{12}CO , ^{13}CO , HCN and HNC, we report on the detection of H_2O and OH emission together with the fine structure lines of [OI] at 63 and 145 μm . The abundance of the latter three species relative to ^{12}CO are $4 \cdot 10^{-2}$, $8 \cdot 10^{-4}$ and 4.5 (approximate value) in the regions where they are produced. We suggest that O-bearing species other than CO are produced in the innermost region of the circumstellar envelope. The UV photons from the central star photodissociate most of the molecular species produced in the AGB phase and allow a chemistry dominated by standard ion-neutral reactions. Not only allow these reactions the formation of O-bearing species, but they also modify the abundances of C-rich molecules like HCN and HNC for which we found an abundance ratio of $\simeq 1$, much lower than in AGB stars. The molecular abundances in the different regions of the circumstellar envelope have been derived from radiative transfer models and our knowledge of its physical structure.

Subject headings: infrared: stars — ISM: molecules — line:
identifications — planetary nebulae: individual (CRL618) — stars:
abundances — stars: carbon

1. Introduction

CRL 618 is one of the few clear examples of an AGB star in the transition phase to the Planetary Nebula stage. It has a compact HII region created by a hot central star (Westbrook 1976, Kwok & Feldman 1981). This object is observed as a bipolar nebula at optical, radio and infrared wavelengths (Carsenty & Solf 1982, Bujarrabal et al. 1988, and Hora et al. 1996). The expansion velocity of the envelope is around 20 km s^{-1} , but CO observations show the presence of a high-velocity outflow with velocities up to 300 km s^{-1} (Cernicharo et al. 1989). High-velocity emission in the $\text{H}_2 v = 1 - 0 \text{ S}(1)$ line is also detected (Burton & Geballe 1986). The high velocity wind and the UV photons from the star perturb the circumstellar envelope (CSE) producing shocks and photodissociation regions (PDRs) which modify the physical and chemical conditions of the gas (Cernicharo et al. 1989, and Neri et al. 1992).

We present in this Letter an ISO (Kessler et al. 1996) Long Wavelength Spectrometer (LWS, Clegg et al. 1996, Swinyard et al. 1996) spectrum between 43 and $197 \mu\text{m}$. In addition to the high-J lines of CO, HCN and HNC, we have discovered, for the first time in a C-rich star, several lines of water vapor, OH and the fine structure lines of [OI]. We discuss the origin of these O-bearing species and we present a model that accounts for the observed emission.

2. Observations and Results

The LWS data were taken on revolution 688. The data have been processed following pipeline number 7. We have used ISAP³ to remove glitches and fringes. The rotational lines (see Fig. 1 and 2) of ^{12}CO (J=14-13 to J=41-40), of ^{13}CO (J=14-13 to J=19-18), HCN, HNC, [OI] (at 63.170 and $145.526 \mu\text{m}$) and C^+ ($^2p_{3/2} - ^2p_{1/2}$ transition at $157.74 \mu\text{m}$) are

observed. Several lines of OH and H₂O are also detected (see Fig. 1 and 2).

The far-infrared spectrum is essentially dominated by the CO lines. All the other species have much lower intensities contrary to what is found in the AGB star IRC+10216 where the HCN lines (from the ground and vibrationally excited states) are as strong as CO. The main difference is due to the physical structure of the CSE of CRL 618 which has a central hole in molecular species filled by a bright HII region. The LWS data also show several weak lines that remain unidentified. We have searched for CH⁺, NH, CH, and other light species. But unlike the case of NGC7027 (Cernicharo et al. 1997) where the PDR has a large spatial size, the PDR of CRL 618 is still close to the central object and has a larger dilution in the LWS beam, which makes the fluxes of these expected species very weak.

The full spectrum of CRL 618 between 2 and 200 μm has been discussed by Cernicharo et al. (1999a et 1999b). In these data, in addition to the species detected in this Letter, they have also reported the presence of several long polyacetylenic chains and the bending modes of HC₃N and HC₅N. However, in the LWS part of the spectrum no pure rotational lines of HC₃N have been found. Nevertheless, the ν_7 band of HC₃N at 44 μm is observed and these data will be published elsewhere. All the observed absorption bands in the mid-infrared arise from gas at $\simeq 250$ K and densities larger than 10^7 cm^{-3} (Cernicharo et al. 1999b).

3. Discussion

Previous works on CRL 618 underlined the presence of a central torus associated with a Photo-Dissociated Region, an extended AGB remnant envelope and high velocity

³The ISO Spectral Analysis Package (ISAP) is a joint development by the LWS and SWS Instrument Teams and Data Centers. Contributing institutes are CESR, IAS, IPAC, MPE, RAL and SRON.

wind regions (HVW) as bipolar lobes (Weintraub et al. 1998, Burton & Geballe 1986, Neri et al. 1992, Carsenty & Solf 1982), whose physical parameters, like the velocity of the gas or the angular size, were investigated (Martin-Pintado et al. 1995, Meixner et al. 1998, Hajian et al. 1996). In order to reproduce the emission, we have to introduce several spatial components, in terms of kinematics, temperature and density based on the knowledge we have of this object. In Fig. 3 we show the adopted geometry: (i) a central torus with the PDR; (ii) the extended AGB remnant envelope; (iii) the lobes (bipolar jets, HVW). We tried to reproduce the LWS observations by simulating a spectrum ($^{12}\text{CO} + ^{13}\text{CO} + \text{HCN} + \text{H}_2\text{O} + \text{OH} + [\text{OI}]$). We first tried to model the ^{12}CO intensities. Then we applied the same parameters to the other molecules allowing to vary only these column densities (see parameters in Table 1). We used a simple LVG-model and obtained satisfactory fits for several molecules (see Fig. 1).

The torus is assumed to be composed of a large neutral shell of $1.1''$ - $1.5''$ size, temperatures from 250 to 800 K, density of $5 \cdot 10^7 \text{ cm}^{-3}$, and an expansion velocity of 20 km s^{-1} . All the detected molecules are present in the 250 K area where most of the CO long wavelength emission comes from. In this region the ^{12}CO column density (N) is 10^{19} cm^{-2} and the ^{13}CO , H_2O , HCN , HNC and OH abundances relative to ^{12}CO are 1/20, 1/25, 1/1000, 1/1000 and 1/1250 respectively. We adopt an ortho-to-para ratio for H_2O around 3. A warmer region in the torus is needed to reproduce the ^{12}CO and ^{13}CO emissions at shorter wavelengths (size $\simeq 0.6''$, $N = 1 \cdot 10^{19} \text{ cm}^{-2}$, $T = 1000 \text{ K}$) and the atomic oxygen emission. As we reach this inner edge of the PDR, only the atomic oxygen seems to be present, whereas all molecular species are photodissociated: $N([\text{OI}]) = 4.5 \cdot 10^{19} \text{ cm}^{-2}$, thus an abundance of 4.5 relative to CO. But this value must be considered as very approximate due to the uncertainties in the size of this region. For the inner radius of the torus, which should be in contact with the HII region, we calculate at a distance of $0.3''$ (10^{16} cm) a UV flux of $20\,000 G_0$ (in units of the local IS radiation flux); this UV illumination is around

200 in the lobes (Latter et al. 1992).

The contribution from the bipolar lobes corresponds to the high velocity gas (200 km s⁻¹, Cernicharo et al. 1989). Previous observations of CO line wings (Cernicharo et al. 1989) and H₂ lines (Herpin et al. in preparation) indicate high velocity and dense shock-heated molecular gas in CRL 618. The contribution of such a region of dense warm shocked gas could dominate the CO high rotational transitions (e.g., Hollenbach & McKee 1989). The high temperature component at high velocity is necessary to produce part of the emission at short wavelengths, since a high temperature component arising from the torus produces too weak intensities. We assume that the bipolar lobes in CO have a size between 1.5'' and 1.7'' with N ranging from $2 \cdot 10^{16}$ to $5 \cdot 10^{18}$ cm⁻², temperatures of 200 K and 1000 K, and density around 10^7 cm⁻³. A contribution from this HVW region for H₂O, OH and [OI] line emission is likely because the molecules in the AGB remnant must be dissociated in the shocks, and in the molecular recombination some O-bearing species could be found like in the inner part of the torus. However, due to the lack of velocity resolution, we can not discriminate both components.

The contribution of the extended AGB remnant envelope is as a colder zone (100 K) for ¹²CO and ¹³CO. The density is around $5 \cdot 10^5$ cm⁻³ and can be compared to what was found by Yamamura et al. (1994) with a mass loss rate of 10^{-4} M_☉/yr for CRL 618, and Neri et al. (1992) ($10^5 - 10^6$ cm⁻³ on the outer edge of the lobes).

For the HVW region and the AGB remnant envelope, we adopt a temperature profile derived from the AGB phase case (Langer & Watson 1984, Herpin 1998), going from 1000 K on the origin of the conical structure (corresponding to the temperature at $6 - 7 \cdot 10^{14}$ cm for a Mira at 3000 K) to 100 K on the exterior, equal to the AGB temperature for dust in the outer part of the envelope (distance $> 10^{16}$ cm) and equal to the value derived by Yamamura et al. (1994).

The model fits well all the CO lines, with small discrepancies at short wavelengths mainly due to the presence of different molecular contributions, to a lower signal-to-noise ratio, and to the fact that some CO lines are mixed with H₂O and OH lines. Nevertheless, all the observed line intensities from J=14-13 up to J=41-40 are reproduced. Our model also fits the observed emissions J=1-0, 2-1 (Cernicharo et al. 1989), 3-2 (Gammie et al. 1989) and 6-5 (unpublished data). A good fit is also obtained for the atomic oxygen lines at 63 μ m and 145 μ m.

Although the ¹³CO emission is weak, the full coverage of the LWS spectrum allows us to derive a [¹²CO / ¹³CO] ratio of 20, lower than the common interstellar value of 40-45, but equal to the detected ratio (20) of Kahane et al. (1992) in this object.

Emission from the HCN wings (Neri et al. 1992) is produced by the strong impact of a primary outflow on a few localized dense clumps in the inner envelope. According to these authors, 70% of the HCN line emission comes from a very compact, spherically symmetric region, and is mainly collisionally excited and strongly affected by self-absorption. They found for the flow (our HVW region) $n(H_2) = \text{few } 10^5 - 10^6 \text{ cm}^{-3}$, $N(\text{CO}) = 2 \cdot 10^{18} \text{ cm}^{-2}$ and $N(\text{HCN}) = \text{few } 10^{17} \text{ cm}^{-2}$, which leads to [HCN/CO] around 10^{-1} . Our high-velocity HCN component (see Table 1) comes from the lobes (HVW) ($N = 2 \cdot 10^{15} \text{ cm}^{-2}$, $T = 1000 \text{ K}$). We find here a ratio [HCN/CO] of 10^{-1} equal to the ratio obtained by Neri et al. (1992). When one considers an abundance of CO of a few 10^{-4} , this leads to a HCN abundance of a few 10^{-5} for our data ([HCN/HNC] $\simeq 10$). This is the expected abundance ratio in a C-rich star in the AGB phase (Cernicharo et al. 1996), and proves that this HCN material has not been processed by the UV photons from the central hot star. Furthermore, Deguchi et al. (1986) found in the disk surrounding the source an abundance of HCN of order of $2 \cdot 10^{-8}$ per H₂ molecule, much less than Neri's value. For this region (see Table 1) we have found a ratio [HCN/CO] of 10^{-3} which leads to an abundance of a few 10^{-7} . We can conclude

that in this region HCN molecules are already efficiently dissociated. The HCN and HNC abundances are similar, which implies a high degree of reprocessing of the molecular gas ($\chi(\text{HCN}) \gg \chi(\text{HNC})$ in standard C-rich AGB stars).

Below 110 μm our models predict that the H_2O and OH lines will be in emission. However, the data seem to indicate that some of these lines could be in absorption or are too weak. Taking into account the reduced sensitivity at these wavelengths and the complex structure of the region, a more detailed model including radiative transfer effects for H_2O and OH, similar to those discussed by Gonzalez-Alfonso & Cernicharo (1999), will need a better knowledge of the physical and geometrical structure of the innermost region of the envelope.

The detection of the fine structure lines of [OI] at 63 and 145 μm gives important insight into the chemical evolution of CRL 618. This indicates that the reservoir of CO molecules formed in the AGB phase of CRL 618 has started to be reprocessed through UV photons and shocks. In CRL 618 Cernicharo et al. (1989) also found H_2CO and Bujarrabal et al. (1988) found HCO^+ . These molecules are not formed efficiently in the AGB phase of a C-rich star. How are they formed in CRL 618 ? As soon as some atomic oxygen is released into the gas phase from the photodissociation of CO, standard ion-neutral reactions can form quickly H_3O^+ and HCO^+ . From the former, and through electronic recombination, H_2O and OH are efficiently produced. We expect these molecules to arise from the innermost region of the envelope where the UV flux is largest. In our models, the column densities in that region are 10^{19} , $5 \cdot 10^{17}$ and $8 \cdot 10^{15} \text{ cm}^{-2}$ for ^{12}CO , H_2O and OH respectively. That means that the abundances of H_2O and OH relative to CO are $4 \cdot 10^{-2}$ and $8 \cdot 10^{-4}$ respectively. These abundances are much lower than those found in the surrounding media of galactic HII regions (Cernicharo et al. 1999a) and suggest that only a small fraction of the UV photons from the central star are reaching the region where

H₂O and OH are formed. In fact, we could expect a dusty region in the torus around the central HII region that protects partially the molecules formed in the AGB phase. Obviously, in the PDR itself all molecules could be destroyed and most of the [OI] dominant coolant in high density PDRs emission is probably coming from that region.

The model derived from the LWS spectra leads to two global regions of molecular emission. One in the torus with several molecular species (¹²CO , ¹³CO , HCN, HNC, H₂O and OH) at moderate temperature. Another in the lobes dominated by CO and HCN molecules. The detected atomic oxygen line is strong and comes from the innermost warm region, the inner edge of the PDR, what proves that CRL 618 is already quite well on its way to the PN stage and that these objects, even in the C-rich case, can produce efficiently O-bearing molecular species.

Acknowledgements We thank spanish DGES and CICYT for funding support for this research under grants PB96-0883 and ESP98-1351E.

REFERENCES

- Bujarrabal V., Gómez-González J., Bachiller R., Martín-Pintado J., 1988, A&A 204, 242
- Burton M.G., Geballe T.R., 1986, MNRAS 223, 13p
- Carsenty U., Solf J., 1982, A&A 106, 307
- Cernicharo J., Guélin M., Martín-Pintado J., Peñalver J., Mauersberger R., 1989, A&A 222, L1
- Cernicharo J. et al., 1996, A&A 315, L201
- Cernicharo J., Liu X.W., González-Alfonso E., Cox P., Barlow M.J., Lim T., Swinyard B.M., 1997, ApJ 483, L65
- Cernicharo J. et al., 1999a, Proceedings of the conference “The Universe as seen by ISO”, ESA SP-427, 655
- Cernicharo J. et al., 1999b, Proceedings of the conference “Astrochemistry Symposium”, Cheju, Korea, August
- Clegg P.E et al., 1996, A&A 315, L38
- Deguchi S., Claussen M.J., Goldsmith P.F., 1986, ApJ 303, 810
- Gammie C.F., Knapp G.R., Young K., Phillips T.G., Falgarone E., 1989, ApJ 345, L87
- Gonzalez-Alfonso, E., Cernicharo J., 1999, ApJ 525, 845
- Hajian A.R., Phillips J. A., Terzian Y., 1996, ApJ 467, 341
- Herpin F., 1998, Ph-D thesis, Université de Bordeaux
- Herpin F. et al., in preparation
- Hollenbach D., McKee C.F., 1989, ApJ 342, 306
- Hora J.L., Deutsch L.K., Hoffmann W.F., Fazio G.G., 1996, AJ 112, 2064

- Kahane C., Cernicharo J., Gómez-González J., Guélin M., 1992, A&A 256, 235
- Kessler M.F. et al., 1996, A&A 315, L27
- Kwok S., Feldman P.A., 1981, ApJ 247, L67
- Langer S.H., Watson W.D., 1984, A&A 284, 751
- Latter W.B et al., 1992, ApJ 389, 347
- Martin-Pintado J., Gaume R. A., Johnston K. J., Bachiller R., 1995, ApJ 446, 687
- Meixner M., Campbell M. T., Welch W. J., Likkell L., 1998, ApJ 509, 392
- Neri R., Garcia-Burillo S., Guélin M., Cernicharo J., Guilloteau S., Lucas R., 1992, A&A 262, 544
- Swinyard B.M. et al.1996, A&A 315, L43
- Weintraub, D. A., Huard, T., Kastner, J.H., Gatley, I., 1998, ApJ 509, 728
- Westbrook W.E., Gezari D.Y., Hauser M.G., Werner M.W., Elias J.H., Neugebauer G., Lo K.Y. 1976, ApJ 209, 94
- Yamamura I., Shibata K.M., Kasuga T., Deguchi S., 1994, ApJ 427, 406

Table 1: Table of the column densities (in cm^{-2}) for all molecules present in each region of the object as depicted in Figure 3 (see text for the adopted parameters).

Molecules	^{12}CO	^{13}CO	HCN	HNC	$\alpha\text{-H}_2\text{O}$	OH	O[I]
TORUS (including PDR)							
T=250 K, $\theta = 1.5''$	10^{19}	$5 \cdot 10^{17}$	10^{16}	10^{16}	$3 \cdot 10^{17}$	$8 \cdot 10^{15}$	
$V = 20 \text{ km s}^{-1}$, $n(\text{H}_2) = 5 \cdot 10^7 \text{ cm}^{-3}$							
T=800 K, $\theta = 1.1''$	$6 \cdot 10^{17}$	$3 \cdot 10^{16}$					
$V = 20 \text{ km s}^{-1}$, $n(\text{H}_2) = 5 \cdot 10^7 \text{ cm}^{-3}$							
T=1000 K, $\theta = 0.6''$	$1 \cdot 10^{19}$	$5 \cdot 10^{17}$					$4.5 \cdot 10^{19}$
$V = 20 \text{ km s}^{-1}$, $n(\text{H}_2) = 7 \cdot 10^7 \text{ cm}^{-3}$							
HVW							
T=200 K, $\theta = 1.7''$	$5 \cdot 10^{18}$	$2.5 \cdot 10^{17}$	$3 \cdot 10^{17}$	$2 \cdot 10^{17}$			
$V = 50 \text{ km s}^{-1}$, $n(\text{H}_2) = 10^7 \text{ cm}^{-3}$							
T=1000 K, $\theta = 1.5''$	$2 \cdot 10^{16}$	$1 \cdot 10^{15}$	$2 \cdot 10^{15}$	10^{14}			
$V = 200 \text{ km s}^{-1}$, $n(\text{H}_2) = 10^7 \text{ cm}^{-3}$							
AGB remnant							
T=100 K, $\theta = 5''$	$7 \cdot 10^{18}$	$3.5 \cdot 10^{17}$	$7 \cdot 10^{17}$	$7 \cdot 10^{15}$			
$V = 20 \text{ km s}^{-1}$, $n(\text{H}_2) = 5 \cdot 10^5 \text{ cm}^{-3}$							

Figure Captions

Fig. 1.— Continuum subtracted spectra of CRL 618 (LW detectors). The result of our model is shown by the continuous line. The lines of CO, ^{13}CO , HCN, H_2O and OH are indicated by arrows while those of HNC are indicated by vertical lines (from J=22-21 at $150.627\ \mu\text{m}$ to J=17-16 at $194.759\ \mu\text{m}$). The C^+ transition is not included in our models; the plot indicates a gaussian fit to this feature.

Fig. 2.— Important detected lines in CRL 618 (continuum subtracted spectra).

Fig. 3.— Simple view of the molecular and emission structure as seen by the model. HVW is for the *High Velocity Wind* region.





